

No C19 Symptoms

Telephone / Video Consult

Offer a F2F appointment if clinically indicated

Tips to deliver good primary care

If your practice has specific reasons why care (eg. blood tests, smears) cannot be delivered due to specific C-19 related risks/capacity issues then consider making good use of the PCAS service or talk to your PCN CD to explore alternatives.

[RCGP/BMA Guidance on workload prioritisation](#)

Staff risk assessment: Ensure the risk/benefit has been considered including a risk assessment of the person carrying out the assessment or procedure using a [recognised health risk assessment tool](#).

Preventative/LTC Care: [See LINK for CCG Guidance](#)

Care Home Visits Checklist <https://www.tamesideandglossopccg.org/clinical>

Caring for vulnerable groups (LCS Bundle):

SMI healthchecks: See [LINK](#) for guidance on CCG expectations.

LD healthchecks: See [LINK](#) for guidance on CCG expectations.

Encouraging optimum self-care

[Signposting patients to self-care resources](#) for optimising health and managing long term conditions.

COVID Vaccination incl complications

Information about local vaccination availability: tameside.gov.uk/covidvaccine

NICE guidance on VITT post-AZ vaccine: [LINK](#)

If patients present following symptoms more than 4 days and within 28 days of AZ vaccine:

- new onset of severe headache, which is getting worse and does not respond to simple painkillers
- an unusual headache which seems worse when lying down or bending over, or may be accompanied by blurred vision, nausea and vomiting, difficulty with speech, weakness, drowsiness or seizures
- new unexplained pinprick bruising or bleeding
- shortness of breath, chest pain, leg swelling or persistent abdominal pain

Direct them to A&E **unless** the person is not acutely unwell, and same day FBC results can be obtained, and if they show thrombocytopenia, the person can be referred to the emergency department immediately.

COVID 19 Testing

Symptomatic staff or patients: www.gov.uk/get-coronavirus-test or 119

Symptomatic staff: Either the same route as symptomatic patients (above) or practice-provided PCR test

Local testing information: tameside.gov.uk/coronavirus/testing

Asymptomatic patient-facing practice staff: Practice-provided lateral flow test (LFT) twice a week and report to <https://www.gov.uk/report-covid19-result>

Asymptomatic members of the public: <https://www.gov.uk/find-covid-19-lateral-flow-test-site>


C19 Symptoms — Cough or fever

(Pts may have myalgia, fatigue, anosmia, sore throat, diarrhoea, congestion or delirium/unexplained deterioration/falls in older people)

Triage Assessment: Phone/Video

This will be done in the first instance by 111/CCAS. However if patients phone their GP surgery then they should be dealt with by the practice and not redirected to 111. CCAS may book directly into GP system via GP Connect.

C19 is the *most likely* cause of symptoms



Alternative diagnosis to C19 more likely (but C19 possible).

Usually no resp symptoms eg. fever due to pyelonephritis, Endocarditis etc

OR

Resp Sx with no fever more likely due to asthma, HF etc

In these circumstances the clinician may decide to risk a brief F2F consultation due to their knowledge of the patient. If this is the case TAKE PRECAUTIONS and use PPE in line with PHE guidance.

Tameside & Glossop CCG/LMC GP Guidance

Vs 28 26/11/2021

Principles for seeing Pts with possible COVID

Consider double triage with colleague.

Person triaging sees the patient.

Restrict building access eg. by entryphone, or allowing 2 people at a time with adequate social distancing.

Consider assessing patients outside.

Clinician wears at least gloves, mask, apron and eye protection. [PPE Guidance](#).

Patient comes in to surgery alone if possible and not to touch anything.

Use the shortest possible path to consulting room and dedicate one room (Red room) in the practice for face to face assessment.

Patient washes hands, and to wear a surgical mask.

Patient brought in for brief exam.

Clean the room surfaces, and equipment with alcohol wipes. Open window(s) to air the room. Remove PPE, wash hands.

Phone patient afterwards to discuss plan and safety net.

Support for GPs, APs and GPNs

Palliative care advice: 24 hour advice line at Willow Wood Hospice, staffed by experienced nurses. 0161 330 5080

Peer GP/PN support phone call from tgccg.gppeersupport@nhs.net Mon-Fri 9-6pm

Check with your PCN resilience lead re. remote O2 satn [Full NHSE Guidance LINK](#)

Videos to help patients to measure their pulse rate and respiratory rate remotely: [Pulse Rate](#) [Respiratory Rate](#)

Supporting patients with post-C19 Symptoms

[GM Support for patients](#)

This link from the BMJ guides GPs/APs in [how to assess patients with possible Post-COVID symptoms](#).

Guidance from BLS/Asthma UK on post-COVID Symptoms [HERE](#).

[Info for patients on symptom management from TGICFT/CCG](#)

On line recovery support <https://www.yourcovidrecovery.nhs.uk/>

T&G OPTIONS: Patients with persistent Sx beyond 12 weeks following COVID or probable COVID can be referred to **TGICFT Post-COVID Syndrome Assessment Clinic**. Referral proforma templates have been sent to Practice Managers to be uploaded into your medical record system.


Updates and Feedback: Please check you are using the most up to date version of this guidance. If any part of the pathway has not worked for you in the way you expect we need to know so that we can sort out problems. If you have any problem or feedback please email tgccg.primarycarereporting@nhs.net

Bronchiolitis Pathway

Clinical Assessment / Management Tool for Children Younger than 1 year old with suspected Bronchiolitis



Management - Primary Care and Community Settings



GMC Best Practice recommends: Record your findings (See "Good Medical Practice" <http://bit.ly/1DPXZ6b>)
**NB: Oximetry is an important part of the assessment and should be measured with an oximeter appropriately designed for infants if available.